arXiv:0803.0235v1 [astro-ph] 3 Mar 2008

Self–enrichment in Globular Clusters: is there a role for the super–asymptotic giant branch stars?

M. L. Pumo[1,2,3], F. D'Antona[1], P. Ventura[1]

[1]*INAF - Osservatorio Astronomico di Roma, via Frascati 33, 00127 Roma, Italy; mlpumo@astrct.oact.inaf.it, dantona@mporzio.astro.it, ventura@mporzio.astro.it*

[2]*Dip.S.F.A. - Università di Palermo, via Archirafi 36, 90123 Palermo, Italy*

[3]*INAF - Osservatorio Astrofisico di Catania, via S. Sofia 78, 95123 Catania, Italy*

Received ... ; accepted ...

ABSTRACT

In four globular clusters (GCs) a non negligible fraction of stars can be interpreted only as a very helium rich population. The evidence comes from the presence of a "blue" main sequence in ω Cen and NGC 2808, and from the the very peculiar horizontal branch morphology in NGC 6441 and NGC 6388. Although a general consensus is emerging on the fact that self–enrichment is a common feature among GCs, the helium content required for these stars is Y $\gtrsim$ 0.35, and it is difficult to understand how it can be produced without any —or, for ω Cen, without a considerable—associated metal enhancement. We examine the possible role of super–AGB stars, and show that they may provide the required high helium. However, the ejecta of the most massive super–AGBs show a global CNO enrichment by a factor of $\simeq$4, due to the dredge–out process occurring at the second dredge up stage. If these clusters show no evidence for this CNO enrichment, we can rule out that at least the most massive super–AGBs evolve into O–Ne white dwarfs and take part in the formation of the second generation stars. This latter hypothesis may help to explain the high number of neutron stars present in GCs. The most massive super–AGBs would in fact evolve into electron–capture supernovae. Their envelopes would be easily ejected out of the cluster, but the remnant neutron stars remain into the clusters, thanks to their small supernova natal kicks.

Subject headings: globular clusters: general — stars: formation — stars: AGB and post-AGB — stars: neutron

1. Introduction

The majority of the inhomogeneities in the chemical composition of Globular Cluster (GC) stars —see e.g. Sneden (1999); Gratton et al. (2004)— appear due to primordial enrichment by hot-CNO cycled material processed in stars belonging to a first stellar generation. Either massive AGB envelopes subject to hot bottom burning, (e.g. Ventura et al. 2001; Ventura, D'Antona, & Mazzitelli 2002) or the envelopes of massive fastly rotating stars (FRMS) (Decressin et al. 2007) are an ideal environment to manufacture elements through nuclear reactions in which proton captures are involved. A second stellar generation would then be born from the ejecta of these stars, either directly, or diluted with pristine material. There are however problems with these scenario. For the AGB progenitors, problems concern the chemistry of the anomalous stars (see Denissenkov & Herwig 2003; Ventura et al. 2004; Fenner et al. 2004, but consider also Ventura & D'Antona 2005), and the requirements for the mass budget necessary to produce a second stellar generation that, today, is generally about as numerous as the first one (D'Antona & Caloi 2004; Bekki & Norris 2006; Prantzos & Charbonnel 2006). The FRMS model is somewhat less problematic concerning the constraints on the IMF, but poses, in any case, severe problems for "normal" GCs, that show a unique metallicity for all stars (with the notable exception of ω Cen): this model requires that the envelopes of these stars contribute to the second star formation stage, but the supernova ejecta are expelled from the clusters and do not enrich the second generation matter by heavy elements.

Both scenarios predict that the stars showing chemical anomalies also must have *enhanced helium abundance*. Helium and the hot–CNO products in the FRMS come to the stellar surface by means of the chemical mixing associated with the transport of angular momentum through the stellar radiative layers, during the phase of hydrogen burning. In the case of AGBs, the helium enrichment is mainly due to the second dredge up (2DU) which precedes the thermal pulse phase. Also the third dredge up (3DU) contributes to helium enrichment in the envelopes, but a further observational constraint is that the sum of CNO abundances remains remarkably constant in normal and anomalous stars (e.g. Cohen & Meléndez 2005). Consequently, the masses which can be responsible for the chemical enrichment must be high enough to be subject to a few episodes of the third dredge up, that brings into the envelope the products of helium burning.

A variation in helium content is immediately reflected on the morphology of the horizontal branch (HB), which amplifies any evolutionary difference among the cluster stars. Helium enrichment by a small factor (up to Y$\sim 0.28-0.30$) allowed in fact an easy interpretation of some puzzles posed by the HB (blue tails, gaps) (D'Antona et al. 2002; D'Antona & Caloi 2004). Helium variations from the MS location are much harder to be detected, but the clustering of $\simeq$20% of ω Cen stars into a "blue" MS (Bedin et al. 2004) was soon interpreted as a MS having an abnormally huge helium content (Y$\sim$0.40) (Norris 2004; Piotto et al. 2005). In contrast to ω Cen, NGC 2808 does not show a metallicity spread (Carretta et al. 2006). Nevertheless, evidence for the presence of a blue MS in this cluster has been found by D'Antona et al. (2005), who interpreted it, again, as built up by stars having Y$\sim$0.40, and were able to correlate them with the bluest side of the HB. Piotto et al. (2007) showed that the MS of this cluster is actually made up by *three* separate MSs, the bluest of which, including 10–15% of stars, must have a helium content Y$\sim 0.35 - 0.40$. The presence of blue MS has not yet been found in other clusters: observationally it is a very difficult problem, and, in addition, this may be a characteristic of the most massive clusters only.

Huge helium overabundances are also needed to explain all the main features of the anomalous HB of the metal rich clusters NGC 6441 and NGC 6388 (Caloi & D'Antona 2007) including: 1) the long extension in luminosity of the red clump; 2) the fact that RR Lyrs have a very long average period, which is unusual for a cluster of high metallicity; 3) the extension into the blue of the HB. This analysis also requires that $\sim$ 15% of the HB population has Y>0.35.

All these observations and their interpretation in terms of helium enrichment must be taken very seriously, but no chemical enrichment mechanism is able to produce the huge amount of helium required for about 15%–20% of the stars in these four clusters without dramatically impacting upon the metal abundance (Karakas et al. 2006; Romano et al. 2007). It is clear that the presence of this extreme population requires some very particular dynamical and chemical conditions to be understood. Nevertheless, the first step is to look for progenitors which may at least provide the required high helium abundance in their ejecta.

The most massive FRMS may be able to eject huge quantities of helium (Decressin et al. 2007). Concerning the AGB stars, Karakas et al. (2006) pointed out that Y$\sim$0.40 is not predicted by AGB models, and this could, in the end, rule out AGB stars as progenitors of the second stellar generation. A scarne Y$\sim$0.29 was the maximum abundance computed from AGB evolution in the models by Ventura et al. (2002) for $5.5M_\odot$, while a single value Y=0.375 is quoted by Karakas et al. (2006) for a $5M_\odot$ model by Campbell et al. (2004). P. Ventura & F. D'Antona (in preparation) have now completed new computation of massive AGBs, whose results revise upward the previous ones (see later). The maximum AGB mass in these recent computation is $6.3M_\odot$, and the possible role of super–AGB stars has not yet been considered. Such models are difficult to be computed, as Carbon burning in semi–degeneracy has to be followed, and the uncertainty in the mass loss rate hampers any conclusions on their final evolution, either towards electron–capture supernova (ecSN) or towards O–Ne white dwarfs.

In the light of recent super–AGB stars models described in Pumo (2006), in this paper we discuss the possible role of these stars for the self–enrichment of GCs. We discuss both the helium abundance and the total CNO abundance versus core mass which can be built up by the most recent AGB and super–AGB models. We find out that the matter lost by the envelopes of super–AGBs may indeed be a good candidate for the high helium population of the four GCs discussed above. However, in the *higher mass tail of the stars developing a O–Ne core* the total CNO is increased by a factor ~ 4. Observations of the super He–rich stars may be able to check this result.

If all the stars in the GCs having a high–He population have standard CNO, as found for all the chemically anomalous stars in many GCs, the most massive super–AGBs must evolve into ecSNe. This conclusion would be of further support to the idea that the presence of large population of neutron stars in GCs implies that they were born with supernova–kicks low enough not to be ejected by the cluster, as expected by the ecSN event (e.g. Ivanova et al. 2007).

2. Super-AGB stellar evolution

AGB stars develop a degenerate carbon–oxygen core and mass loss prevents them to reach the mass limit for degenerate explosive carbon ignition. Super–AGBs are defined as stars which ignite carbon in conditions of semi–degeneracy, thus non explosively, but are not able to ignite hydrostatic Neon burning in the resulting O–Ne core. Consequently, degeneracy increases in the core and these stars may undergo thermal pulses (e.g. Iben et al. 1997; Ritossa et al. 1999; Siess 2006) and lose mass as "normal" (but quite massive and luminous) AGB stars. Depending on the competition between the mass loss rate and the core growth, they may then evolve into massive O–Ne white dwarfs (Nomoto 1984) or as ecSN, electrons being captured first by ^{24}Mg and ^{24}Na then by ^{20}Ne and ^{20}F nuclei when the core mass reaches the Chandrasekhar mass. The 2DU occurs also in super–AGBs and, as in the massive AGBs, it reduces the mass of the H–exhausted core (e.g. Pumo & Siess 2006), while the helium abundance increases in the external hydrogen envelope (e.g. Siess & Pumo 2006). If mass loss is strong enough to avoid the super–AGB evolving to ecSN, as soon as super–AGBs evolve in the GCs, the supernova epoch is finished, and we may hypothesize that the second phase of star formation begins from the super–AGB ejecta.

3. Helium and CNO in massive AGBs and super–AGBs

As a function of the Carbon Oxygen core mass M_c, we report in Figure 1 the helium content in the AGB ejecta of metallicity Z=10^{-3}, for the models from 4 to 6.3$M_\odot$ by P. Ventura & F. D'Antona (in preparation) including α–elements enhancement in the opacities, and core overshooting. An important difference from Ventura et al. (2002) models is the following: in the present models, also overshooting at the bottom of the convective envelope is included, adopting the same parameter which describes core overshooting. This latter is calibrated to account for the width of the MS of intermediate mass stars. The effect of the envelope overshooting is to enhance the extent of the 2DU: fixed the initial mass, the remnant core mass at the beginning of the AGB evolution is then *smaller*, and the resulting helium content in the envelope is *larger*, as can be seen by comparing the values of Figure 1 with Table 1 in Ventura et al. (2002).

We also plot the helium content at the 2DU for the super–AGB models of metallicity Z=10^{-3} from 7.5$M_\odot$ to 9.5$M_\odot$ (Pumo 2006) computed without overshooting. In this case M_c is the C–O mass after the 2DU. M_c is indeed the important physical parameter which determines the final fate of the star. The initial mass difference between the AGB models with overshooting and super–AGB models without overshooting is, for the same core mass, $\sim$1$M_\odot$. For super–AGBs, we do not have the helium mass fraction of the ejecta, but these stars do not go through efficient episodes of 3DU, so the plotted value is a good approximation. As we see, the super–AGB models reach a helium content in the ejecta larger than the standard AGBs, and well approaching the values needed to be consistent with the super–He rich stars in GCs. The new AGB models and the super–AGB models shown here, computed by a different program and input physics (Siess 2006; Siess & Pumo 2006) are in good agreement with each other, at the same core mass.

From Figure 1 we see that indeed the super–AGBs could be the progenitors of the very high helium population found in the most massive GCs. For a Salpeter's initial mass function, the mass budget in the ejecta from 6.5$M_\odot$(the minimum mass for super–AGB evolution, considering models with overshooting) to 8$M_\odot$(see later) is about 50% of the mass budget contained in the ejecta of normal AGBs from 5$M_\odot$to 6.5$M_\odot$. Therefore, from the super–AGBs mass budget, a 15% fraction of very high helium stars can be born, with the same mechanism which can give origin to another $\sim$30% of moderately helium rich stars, as in the cluster NGC 2808 (D'Antona et al. 2005). How these stars can form directly from the super–AGB ejecta, so that their helium abundance remains as large as the required Y $\gtrsim$ 0.35, is a different problem.

Figure 1 also shows the total CNO enrichment F$_{CNO}$, the sum of CNO abundances with respect to their initial sum, in the ejecta of AGBs and at the 2DU for the super–AGBs. The

AGB enrichment is larger than ∼2 for $M_c \leq 0.9M_{\odot}$, that is for M$\leq 5M_{\odot}$. As we have noticed, the sum of CNO abundances apparently is constant (within a factor 2) in GC stars. On this basis, we should conclude that stars of M$<5M_{\odot}$ should not contribute to the second stellar generation –unless their ejecta are diluted with pristine matter, so that the constancy of CNO is preserved. For the super–AGBs, we also find a sharp increase of F$_{CNO}$, reaching a factor 4 for the largest mass in exam. This is due to the occurrence of the so called dredge–out process in this model (Iben et al. 1997; Ritossa et al. 1999). The helium mass fraction and the overall CNO content are the most robust predictions from these theoretical computations, due to the large uncertainties associated to both the extension of the 3DU and the strength of hot bottom burning (HBB) within super–AGB models (e.g. Izzard & Poelarends 2006; Pumo & Siess 2006). We also comment the individual abundances providing F$_{CNO}$. The CNO abundances given for $M \leq 6.3M_{\odot}$are the average mass fractions of the ejecta. As HBB is very efficient, the values provide very small carbon ($X_{^{12}C} \simeq 1.1 \cdot 10^{-4}$ for 4-5$M_{\odot}$models down to $X_{^{12}C} \simeq 1.6 \cdot 10^{-5}$ at $6.3M_{\odot}$), and low oxygen (from $X_{Ø16} \simeq 6.3 \cdot 10^{-4}$ at $4M_{\odot}$to $X_{Ø16} \simeq 10^{-4}$ at $6.3M_{\odot}$); nitrogen is very high, especially in the $4M_{\odot}$, where the effect of the 3DU is more visible ($X_{^{14}N} \simeq 2.6 \cdot 10^{-3}$). On the contrary, for the super–AGBs, we consider the values at the 2DU. The carbon abundance increases from $X_{^{12}C} \simeq 8.9 \cdot 10^{-5}$ for the $7.5M_{\odot}$model to $X_{^{12}C} \simeq 8.1 \cdot 10^{-4}$ at $9M_{\odot}$. The nitrogen is about constant ($X_{^{14}N} \simeq 2.4 \cdot 10^{-4}$), whereas the oxygen abundance is constant ($X_{Ø16} \simeq 3.8 \cdot 10^{-4}$) up to $8.5M_{\odot}$, and increases up to $X_{Ø16} \simeq 4.4 \cdot 10^{-4}$ for the $9M_{\odot}$model: this is due to the deeper sinking of the 2DU when increasing the mass. In the $9.5M_{\odot}$model, the dredge–out process rises $X_{^{12}C}$ to $\simeq 1.8 \cdot 10^{-3}$ and $X_{Ø16}$ to $\simeq 7.7 \cdot 10^{-4}$. The following evolution will reduce oxygen and carbon and increase nitrogen, if HBB is efficient in these stars as it is in the massive AGBs (this depends on the convective model adopted; see Ventura & D'Antona 2005). As for the 3DU, there are different and incomplete literature results (see Izzard & Poelarends 2006, for summary). In any case, it can only act to increase the values of total CNO abundances given in Figure 1.

How many stars would be affected if all the super–AGBs contribute to the second stellar generation? If all the super–AGBs contribute to the very helium rich population, the fraction of high–CNO stars, assuming a Salpeter's IMF, would be ≃10%! This is a small fraction of the cluster stars, if we consider that the super–He rich stars are 15–20% of the total population, but spectroscopic observations of a large sample of stars could falsify this hypothesis.

4. Do the most massive super–AGBs explode?

Should the constancy of CNO be observationally confirmed also for the most helium rich population showing up in the quoted 4 clusters, we will be able to conclude that the most massive super–AGBs did not take part in the process of forming the second stellar generation in GCs. This may mean: 1) that there were reasons, independent from the super–AGB evolution, by which the self–enrichment process was effective either only at earlier ages (due to the envelopes of massive stars) or at a later age (during the normal, massive AGB evolution) 2) or that the self–enrichment was forbidden by the injection in the cluster gas of the energy due to the ecSN explosion of the stars which undergo the dredge out. Although the explosion energy of such event is significantly lower than inferred for core collapse SN (Dessart et al. 2006), it is probably more than sufficient to expel the SN ejecta from most clusters. The conclusion then would be that at least a fraction of super–AGBs must evolve into ecSN, reinforcing the idea that ecSN channel for super–AGB stars could occur (see e.g. Siess & Pumo 2006; Pumo & Siess 2006; Poelarends et al. 2007).

5. Neutron stars in Globular Clusters

The final fate as ecSN of —at least a fraction— of super–AGBs is also required by the recent suggestion that NS in GCs should be mainly formed by the ecSN channel.

The high number of neutron stars (NS) in GCs, whose presence is revealed mainly by the very high number of millisecond pulsars, remains a puzzle. The massive GC 47Tuc was predicted to contain more than 1000 NS (see e.g. Pfahl et al. 2002). Now this figure can perhaps be reduced to ∼300–600 (Heinke et al. 2005). The velocity distribution of young pulsars in the Galaxy shows that, at formation, the NS receives a "natal kick", most likely due to the asymmetry in the supernova ejecta Fryer (2004). Pfahl et al. (2002) have shown that only up to 8% of NS can be retained in the clusters. The kick velocity distribution which they used (Hansen & Phinney 1997) has been recently updated by Hobbs et al. (2005), who find a higher mean velocity and velocity dispersion. Using these new data, simulations of the retention factor by Ivanova et al. (2007) show that almost no NS can be retained for an escape velocity of 40Km$\cdot$s^{-1}. A natural way to solve this problem, is to hypothesize that there is a stellar population which evolves into ecSN, that probably have *much smaller* natal kicks (Ivanova et al. 2007), possibly proportional to their reduced explosion energy. This is also suggested by the existence of a class of massive X–ray binaries with small eccentricities. van den Heuvel (2007) also supports the evidence of two channels for NS formation —with either small or large natal kicks— from an exam of the double neutron star binaries.

We are grateful to R. G. Izzard for his helpful discussions.

REFERENCES

Bedin, L. R., Piotto, G., Anderson, J., Cassisi, S., King, I. R., Momany, Y., Carraro, G. 2004, ApJ, 605, L125

Bekki, K., & Norris, J. E. 2006, ApJ, 637, L109

Caloi, V. & D'Antona, F. 2007, A&A, 463, 949

Campbell, S. W., Fenner, Y., Karakas, A. I., Lattanzio, J. C., & Gibson, B. K. 2004, Memorie della Societa Astronomica Italiana, 75, 735

Carretta, E., Bragaglia, A., Gratton, R. G., Leone, F., Recio-Blanco, A., & Lucatello, S. 2006, A&A, 450, 523

Cohen, J. G., & Meléndez, J. 2005, AJ, 129, 303

D'Antona, F., Caloi, V., Montalbán, J., Ventura, P., & Gratton, R. 2002, A&A, 395, 69

D'Antona, F., & Caloi, V. 2004, ApJ, 611, 871

D'Antona, F., Bellazzini, M., Caloi, V., Fusi Pecci, F., Galleti, S., & Rood, R. T. 2005, ApJ, 631, 868

Decressin, T., Meynet, G., Charbonnel, C., Prantzos, N., & Ekström, S. 2007, A&A, 464, 1029

Denissenkov, P. A., & Herwig, F. 2003, ApJ, 590, L99

Dessart, L., Burrows, A., Ott, C. D., Livne, E., Yoon, S.-C., & Langer, N. 2006, ApJ, 644, 1063

Fenner,Y., Campbell, S., Karakas, A. I., Lattanzio, J. C., & Gibson, B. K. 2004, MNRAS, 353, 789

Fryer, C. L. 2004, ApJ, 601, L175

Gratton, R., Sneden, C., & Carretta, E. 2004, ARA&A, 42, 385

Hansen, B. M. S., & Phinney, E. S. 1997, MNRAS, 291, 569

Heinke, C. O., Grindlay, J. E., Edmonds, P. D., Cohn, H. N., Lugger, P. M., Camilo, F., Bogdanov, S., & Freire, P. C. 2005, in AIP Conf. Proc., Interacting Binaries: Accretion, Evolution, and Outcomes, 797, 40

Hobbs, G., Lorimer, D. R., Lyne, A. G., & Kramer, M. 2005, MNRAS, 360, 974

Iben, I. J., Ritossa, C., & Garcia-Berro, E. 1997, ApJ, 489, 772

Ivanova, N., Heinke, C., Rasio, F. A., Belczynski, K., & Fregeau, J. 2007, preprint (arXiv:0706.4096v1)

Izzard, R. G., & Poelarends, A. J. T. 2006, Memorie della Societa Astronomica Italiana, 77, 840

Karakas, A. I., Fenner, Y., Sills, A., Campbell, S. W., & Lattanzio, J. C. 2006, ApJ, 652, 1240

Lattanzio, J., Karakas, A., Campbell, S., Elliott, L., & Chieffi, A. 2004, Memorie della Societa Astronomica Italiana, 75, 322

Nomoto, K. 1984, ApJ, 277, 791

Norris, J. E. 2004, ApJ, 612, L25

Pfahl, E., Rappaport, S., & Podsiadlowski, P. 2002, ApJ, 573, 283

Piotto, G., et al. 2005, ApJ, 621, 777

Piotto, G., et al. 2007, ApJ, 661, L53

Poelarends, A. J. T., Herwig, F., Langer, N., Heger, A. 2007, preprint (arXiv:0705.4643v1)

Prantzos, N., & Charbonnel, C. 2006, A&A, 458, 135

Pumo, M.L. 2006, PhD thesis, University of Catania (Italy)

Pumo, M. L., & Siess, L. 2006, in ASP Conf. Ser., Why galaxies care about AGB stars, ed. F. Kerschbaum, C. Charbonnel, B. Wing, in press

Ritossa, C., García-Berro, E., & Iben, I. J. 1999, ApJ, 515, 381

Romano, D., Matteucci, F., Tosi, M., Pancino, E., Bellazzini, M., Ferraro, F. R., Limongi, M., & Sollima, A. 2007, MNRAS, 376, 405

Siess, L. 2006, A&A, 448, 717

Siess, L., & Pumo, M. L. 2006, Memorie della Societa Astronomica Italiana, 77, 822

Sneden C. 1999, Ap&SS, 265, 145

van den Heuvel, E. P. J. 2007, American Institute of Physics Conference Series, 924, 598

Ventura, P., D'Antona, F., Mazzitelli, I., & Gratton, R. 2001, ApJ, 550, L65

Ventura, P., D'Antona, F., & Mazzitelli, I. 2002, A&A, 393, 215

Ventura, P., & D'Antona, F. 2005, A&A, 431, 279

Ventura, P., D'Antona, F., & Mazzitelli, I. 2004, Memorie della Societa Astronomica Italiana, 75, 335

This preprint was prepared with the AAS LaTeX macros v5.2.



Fig. 1.— The triangles plus dashed line show the helium content in the ejecta of AGB stars as function of the C–O core mass, in the computations by P. Ventura & F. D'Antona (in preparation). The initial masses are, from left to right, 4, 4.5, 5, 5.5, 6, 6.3$M_\odot$. The triangles plus continuous line show the helium abundance in the envelope after the 2DU in the super–AGBs by Pumo (2006). Masses from left to right are 7.5, 8, 8.5, 9, 9.5$M_\odot$. The circles show the ratio of total CNO abundance (F_{CNO}) with respect to the initial value for the same models.